

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 921 194 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

09.06.1999 Bulletin 1999/23

(51) Int Cl.⁶ **C12N 15/12, A61K 38/17,**

G01N 33/68, C12N 1/21,

C07K 14/705, A61K 38/19,

C07K 14/52, C12N 15/19

(21) Application number: **98309632.2**

(22) Date of filing: **25.11.1998**

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

Designated Extension States:

AL LT LV MK RO SI

(72) Inventor: **Song, Ho Yeong**

Carmel, Indiana 46032 (US)

(74) Representative: **Denholm, Anna Marie et al**

Eli Lilly and Company Limited,

Lilly Research Center,

Erl Wood Manor

Windlesham, Surrey GU20 6PH (GB)

(30) Priority: **26.11.1997 US 66577 P**

11.08.1998 US 96173 P

(71) Applicant: **ELI LILLY AND COMPANY**

Indianapolis, Indiana 46285 (US)

(54) **TRAILLK-2: a member of the TNF ligand family**

(57) The invention provides isolated nucleic acid compounds, proteins and fragments thereof, said proteins being related to the family of TNF ligands. Also provided are vectors and transformed heterologous host

cells for expressing the protein and a method for identifying compounds that bind and/or modulate the activity of said proteins, and a method for treating cancerous growths in a patient in need thereof.

EP 0 921 194 A2

Description

[0001] This application claims the benefit of U.S. Provisional Application No. 60/066,577, filed November 26, 1997, and U.S. Provisional Application No. 60/096,173, filed August 11, 1998.

[0002] This invention relates to recombinant DNA technology. In particular the invention pertains to a TNF ligand family gene, and its encoded protein. Also contemplated are methods for identifying compounds that bind said ligand, and methods for treating oncological disease in mammals.

[0003] Apoptosis (*i.e.* programmed cell death), is a process fundamental to the normal development and homeostasis of multicellular organisms. Deregulation of programmed cell death leads to a number of human diseases, including cancer, neurodegenerative disorders, and acquired immunodeficiency syndrome. The cell death machinery comprises effectors, activators, and negative regulators. Certain cytokines of the tumor necrosis factor (TNF) ligand family and their cognate receptors, including TNFR-1 and Fas (also known as Apo-1 or CD95), are classic triggers of the suicide response. TNF is the prototypic member of an emerging family of cytokines that function as prominent mediators of immune regulation and the inflammatory response. Eight other members of the TNF family are known, including lymphotoxin (LT α , TNF β), lymphotoxin β (LT β), and ligands for CD40, CD30, CD27, OX40, 4-1BB, and Fas (APO-1). With one exception, all TNF ligands are type II membrane proteins, having homology at the C-terminal end. The exception, LT α , appears to be a secreted protein that also exists as a cell surface-associated member, termed LT β . In addition, a proteolytically processed soluble form of TNF has been studied.

[0004] The TNF ligands interact with a parallel family of some twelve homologous receptors, characterized by cysteine-rich pseudorepeats in the extracellular region. As with the TNF ligands, the TNF receptors are variably expressed in a variety of cell types, including B cells, T cells, dendritic cells, and macrophages.

[0005] TNF and Fas ligand (also known as Apo-1L or CD95L) induce apoptosis by binding to receptors TNFR-1 and Fas. These receptors contain a domain (termed "death domain") that mediates the assembly of a signaling complex, leading to the recruitment of pro-apoptotic proteases.

[0006] Another member of the TNF ligand family, termed TRAIL (also known as Apo-2L), has been identified. Like the Fas ligand (FasL), TRAIL induces rapid apoptosis in transformed cell lines of diverse origin. Unlike FasL, whose transcripts are predominantly restricted to stimulated T cells, TRAIL expression is detected in many normal human tissues. This suggests that TRAIL is a member of the TNF ligand family that has marked pro-apoptotic potential for transformed cells. However, the inability of TRAIL to bind TNFR-1, Fas, or the recently identified death domain-containing receptor DR3 (also called Ws1-1, Apo-3, and TRAMP) suggests that TRAIL may interact with a yet unknown member of the TNF-receptor family.

[0007] The present invention provides isolated nucleic acid molecules and novel members of the TNF ligand family, termed herein "TRAILLK-2". Having the cloned TRAILLK-2 gene enables the production of recombinant protein, the isolation of orthologous genes from other organisms, and/or paralogous genes from the same organism, chromosome mapping studies, and the implementation of large scale screens to identify compounds that bind said receptor protein, as a means to identify potential pharmaceutical compounds for modulating the biological activity thereof. The proteins and peptides described herein are also useful therapeutic agents *per se* and for developing new compounds for treating cancer, and, among other things, as feed additives.

[0008] In one embodiment the present invention relates to an isolated nucleic acid molecule encoding TRAILLK-2 protein.

[0009] In another embodiment, the invention relates to a nucleic acid molecule comprising the nucleotide sequence identified as SEQ ID NO:1.

[0010] In another embodiment, the present invention relates to a nucleic acid that encodes SEQ ID NO:2, or fragment thereof.

[0011] In another embodiment, the present invention relates to a nucleic acid that is at least 75% identical to a nucleic acid that encodes SEQ ID NO:2, or fragment thereof.

[0012] In another embodiment, the present invention relates to a nucleic acid that hybridizes to SEQ ID NO:1 under high stringency conditions and encodes a protein that is capable of inducing apoptosis, and/or in treating cancer, and/or in preventing or inhibiting cancerous tumor growth, and/or in causing shrinkage of a cancerous tumor.

[0013] In another embodiment, the present invention relates to a nucleic acid that hybridizes to SEQ ID NO:1 under low stringency conditions and encodes a protein that is capable of inducing apoptosis and/or in treating cancer, and/or in preventing or inhibiting cancerous tumor growth, and/or in causing shrinkage of a cancerous tumor.

[0014] In another embodiment the present invention relates to an isolated protein molecule, or functional fragment thereof, wherein said protein molecule comprises the sequence identified as SEQ ID NO:2.

[0015] In yet another embodiment, the present invention relates to a recombinant DNA vector that incorporates the TRAILLK-2 gene (SEQ ID NO:1) in operable-linkage to gene expression sequences, enabling the gene to be transcribed and translated in a host cell.

[0016] In still another embodiment the present invention relates to host cells that have been transformed or trans-

fectured with the cloned TRAILLK-2 gene such that said gene is expressed in the host cell.

[0017] This invention also provides a method of determining whether a nucleic acid sequence of the present invention, or fragment thereof, is present within a nucleic acid-containing sample, comprising contacting the sample under suitable hybridization conditions with a nucleic acid probe of the present invention.

[0018] In a still further embodiment, the present invention relates to a method for treating cancer, and/or in preventing or inhibiting cancerous tumor growth, and/or in causing shrinkage of a cancerous tumor.

Definitions

[0019] "Apoptosis" refers to the phenomenon of programmed cell death that is fundamental to the normal development and homeostasis of multicellular organisms. Deregulation of apoptosis leads to a number of human diseases, including cancer, neurodegenerative disorders, and AIDS. Certain cytokines of the TNF ligand family are triggers of apoptosis.

[0020] The terms "complementary" or "complementarity" as used herein refer to the capacity of purine and pyrimidine nucleotides to associate through hydrogen bonding to form double stranded nucleic acid molecules. The following base pairs are related by complementarity: guanine and cytosine; adenine and thymine; and adenine and uracil. As used herein, "complementary" means that the aforementioned relationship applies to substantially all base pairs comprising two single-stranded nucleic acid molecules over the entire length of said molecules. "Partially complementary" refers to the aforementioned relationship in which one of two single-stranded nucleic acid molecules is shorter in length than the other such that a portion of one of the molecules remains single-stranded.

[0021] "Conservative substitution" or "conservative amino acid substitution" refers to a replacement of one or more amino acid residue(s) in a protein or peptide as stipulated in Table 1.

[0022] "Fragment thereof" refers to a fragment, piece, or sub-region of a nucleic acid or protein molecule whose sequence is disclosed herein, such that the fragment comprises 5 or more amino acids, or 10 or more nucleotides that are contiguous in the parent protein or nucleic acid molecule. Fragment thereof may or may not retain biological activity. For example, a fragment of a protein disclosed herein could be used as an antigen to raise a specific antibody against the parent protein molecule. When referring to a nucleic acid molecule, "fragment thereof" refers to 10 or more contiguous nucleotides, derived from the parent nucleic acid, and also, owing to the genetic code, to the complementary sequence. For example if the fragment entails the sequence 5'-AGCTAG-3', then "fragment thereof" would also include the complementary sequence, 3'-TCGATC-5'.

[0023] The term "fusion protein" denotes a hybrid protein molecule not found in nature comprising a translational fusion or enzymatic fusion in which two or more different proteins or fragments thereof are covalently linked on a single polypeptide chain.

[0024] "Functional fragment" or "functionally equivalent fragment", as used herein, refers to a region, or fragment of a full length protein, or sequence of amino acids that, for example, comprises an active site, or any other conserved motif, relating to biological function. Functional fragments are capable of providing a substantially similar biological activity as a full length protein disclosed herein, *in vivo* or *in vitro*, viz. the capacity to promote apoptosis. Functional fragments may be produced by cloning technology, or as the natural products of alternative splicing mechanisms.

[0025] "Host cell" refers to any eucaryotic or procaryotic cell that is suitable for propagating and/or expressing a cloned gene contained on a vector that is introduced into said host cell by, for example, transformation or transfection, or the like.

[0026] TRAILLK-2 refers to a gene (SEQ ID NO:1) and a protein (SEQ ID NO:2) or peptide encoded thereby. TRAILLK-2 is a member of the family of TNF ligands. This family mediates a variety of biological effects including apoptosis, especially in tumor cells, and induction of the immune response.

[0027] The term "homolog" or "homologous" describes the relationship between different nucleic acid molecules or amino acid sequences in which said sequences or molecules are related by partial identity or similarity at one or more blocks or regions within said molecules or sequences.

[0028] The term "hybridization" as used herein refers to a process in which a single-stranded nucleic acid molecule joins with a complementary strand through nucleotide base pairing. "Selective hybridization" refers to hybridization under conditions of high stringency. The degree of hybridization depends upon, for example, the degree of homology, the stringency of hybridization, and the length of hybridizing strands.

[0029] "Isolated nucleic acid compound" refers to any RNA or DNA sequence, however constructed or synthesized, which is locationally distinct from its natural location.

[0030] A "nucleic acid probe" or "probe" as used herein is a labeled nucleic acid compound which hybridizes with another nucleic acid compound. "Nucleic acid probe" means a single stranded nucleic acid sequence that will combine with a complementary or partially complementary single stranded target nucleic acid sequence to form a double-stranded molecule. A nucleic acid probe may be an oligonucleotide or a nucleotide polymer. A probe will usually contain a detectable moiety which may be attached to the end(s) of the probe or be internal to the sequence of the probe.

[0031] The term "orthologue" or "orthologous" refers to two or more genes or proteins from different organisms that exhibit sequence homology.

[0032] The term "paralogue" or "paralogous" refers to two or more genes or proteins within a single organism that exhibit sequence homology.

[0033] The term "plasmid" refers to an extrachromosomal genetic element. The plasmids disclosed herein are commercially available, publicly available on an unrestricted basis, or can be constructed from readily available plasmids in accordance with published procedures.

[0034] A "primer" is a nucleic acid fragment which functions as an initiating substrate for enzymatic or synthetic elongation of, for example, a nucleic acid molecule.

[0035] The term "promoter" refers to a nucleic acid sequence that directs transcription, for example, of DNA to RNA. An inducible promoter is one that is regulatable by environmental signals, such as carbon source, heat, or metal ions, for example. A constitutive promoter generally operates at a constant level and is not regulatable.

[0036] "Recombinant DNA cloning vector" as used herein refers to any autonomously replicating agent, including, but not limited to, plasmids and phages, comprising a DNA molecule to which one or more additional DNA segments can or have been incorporated.

[0037] The term "recombinant DNA expression vector" or "expression vector" as used herein refers to any recombinant DNA cloning vector, for example a plasmid or phage, in which a promoter and other regulatory elements are present thereby enabling transcription of an inserted DNA, which may encode a protein.

[0038] The term "stringency" refers to hybridization conditions. High stringency conditions disfavor non-homologous base pairing. Low stringency conditions have the opposite effect. Stringency may be altered, for example, by temperature and salt concentration.

[0039] "Low stringency" conditions comprise, for example, a temperature of about 37° C or less, a formamide concentration of less than about 50%, and a moderate to low salt (SSC) concentration; or, alternatively, a temperature of about 50° C or less, and a moderate to high salt (SSPE) concentration, for example 1M NaCl.

[0040] "High stringency" conditions comprise, for example, a temperature of about 42° C or less, a formamide concentration of less than about 20%, and a low salt (SSC) concentration; or, alternatively, a temperature of about 65° C, or less, and a low salt (SSPE) concentration. For example, high stringency conditions comprise hybridization in 0.5 M NaHPO₄, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65° C (Ausubel, F.M. et al. Current Protocols in Molecular Biology, Vol. I, 1989; Green Inc. New York, at 2.10.3).

[0041] The symbol "N" in a nucleic acid sequence refers to adenine ("A"), guanine ("G"), cytosine ("C"), thymine ("T"), or uracil ("U").

[0042] "SSC" comprises a hybridization and wash solution. A stock 20X SSC solution contains 3M sodium chloride, 0.3M sodium citrate, pH 7.0.

[0043] "SSPE" comprises a hybridization and wash solution. A 1X SSPE solution contains 180 mM NaCl, 9mM Na₂HPO₄, 0.9 mM NaH₂PO₄ and 1 mM EDTA, pH 7.4.

[0044] "Substantially pure," used in reference to a peptide or protein, means separation from other cellular and non-cellular molecules, including other protein molecules. A substantially pure preparation would be about at least 85% pure; preferably about at least 95% pure. A "substantially pure" protein can be prepared by a variety of techniques, well known to the skilled artisan, including, for example, the IMAC protein purification method.

[0045] "Treating" as used herein describes the management and care of a patient for the purpose of combating the disease, condition, or disorder and includes the administration of a protein of the present invention to prevent the onset of the symptoms or complications, alleviating the symptoms or complications, or eliminating the disease, condition, or disorder. Treating as used herein includes the administration of the protein for cosmetic purposes. A cosmetic purpose seeks to control, for example, the weight of a mammal to improve bodily appearance.

[0046] The term "vector" as used herein refers to a nucleic acid compound used for introducing exogenous or endogenous DNA into host cells. A vector comprises a nucleotide sequence which may encode one or more protein molecules. Plasmids, cosmids, viruses, and bacteriophages, in the natural state or which have undergone recombinant engineering, are examples of commonly used vectors.

[0047] The various restriction enzymes disclosed and described herein are commercially available and the manner of use of said enzymes including reaction conditions, cofactors, and other requirements for activity are well known to one of ordinary skill in the art. Reaction conditions for particular enzymes were carried out according to the manufacturer's recommendation.

[0048] The TRAILLK-2 gene encodes a novel, membrane-bound protein that is related to the TNF family. The TRAILLK-2 cDNA comprises a DNA sequence specified herein by SEQ ID NO:1. The coding region begins at base pair 49 of SEQ ID NO:1 and extends through base pair 798 of SEQ ID NO:1. Those skilled in the art will recognize that owing to the degeneracy of the genetic code, numerous "silent" substitutions of nucleotide base pairs could be introduced into the sequence identified as SEQ ID NO:1 without altering the identity of the encoded amino acid(s) or protein product. All such substitutions are intended to be within the scope of the invention.

[0049] Also contemplated by the present invention are related proteins and related functional fragments such as, for example, smaller alternatively spliced forms. Related proteins comprise a genus in which amino acid substitutions of the primary sequence disclosed in SEQ ID NO:2 is altered by substitution, or replacement, or deletion, or insertion at one or more amino acid positions, such that TNF ligand function is maintained.

[0050] Functional fragments are conveniently identified as fragments of an intact TRAILLK-2 protein that retain the capacity to induce apoptosis.

[0051] Amino acid substitution modifications can be made in accordance with the following Table.

ORIGINAL RESIDUE	EXEMPLARY SUBSTITUTIONS
ALA	SER
ARG	LYS
ASN	GLN; HIS
ASP	GLU
CYS	SER
GLN	ASN
GLU	ASP
GLY	PRO
HIS	ASN, GLN
ILE	LEU, VAL
LEU	ILE, VAL
LYS	ARG, GLN, GLU
MET	LEU, ILE
PHE	MET, LEU, TYR
SER	THR
THR	SER
TRP	TYR
TYR	TRP, PHE
VAL	ILE, LEU

Fragments of proteins

[0052] One embodiment of the instant invention provides fragments of the proteins disclosed that may or may not be biologically active. Such fragments are useful, for example, as an antigen for producing an antibody to said proteins.

[0053] Fragments of the proteins disclosed herein may be generated by any number of suitable techniques, including chemical synthesis of any portion of SEQ ID NO:2, proteolytic digestion of SEQ ID NO:2, or most preferably, by recombinant DNA mutagenesis techniques, well known to the skilled artisan. See, e.g. K. Struhl, "Reverse biochemistry: Methods and applications for synthesizing yeast proteins *in vitro*," *Meth. Enzymol.* 194, 520-535. For example, in a preferred method, a nested set of deletion mutations are introduced into the intact gene (SEQ ID NO:1) encoding TRAILLK-2 such that varying amounts of the protein coding region are deleted, either from the amino terminal end, or from the carboxyl end of the protein molecule. This method can also be used to create internal fragments of the intact protein in which both the carboxyl and amino terminal ends are removed. Several appropriate nucleases can be used to create such deletions, for example *BaB1*, or in the case of a single stranded nucleic acid molecule, mung bean nuclease. For simplicity, it is preferred that the TRAILLK-2 gene be cloned into a single-stranded cloning vector, such as bacteriophage M13, or equivalent. If desired, the resulting gene deletion fragments can be subcloned into any suitable vector for propagation and expression of said fragments in any suitable host cell.

[0054] The present invention also provides fragments of the proteins disclosed herein wherein said fragments retain receptor activity. As used herein, "functional fragments" refer to fragments of SEQ ID NO:2 that retain and exhibit, under appropriate conditions, measurable biological activity, for example, the capacity to induce apoptosis.

[0055] Functional fragments of the proteins disclosed herein may be produced as described above, preferably using cloning techniques to engineer smaller versions of the intact gene, lacking sequence from the 5' end, the 3' end, from both ends, or from an internal site. Fragments may be tested for biological activity using any suitable assay, for example, the ability of a protein fragment to induce apoptosis, *in vivo* or *in vitro*.

Gene Isolation Procedures

[0056] Those skilled in the art will recognize that the TRAILLK-2 gene could be obtained by a plurality of recombinant DNA techniques including, for example, hybridization, polymerase chain reaction (PCR) amplification, or de novo DNA synthesis. (See e.g., T. Maniatis *et al.* Molecular Cloning: A Laboratory Manual, 2d Ed. Chap. 14 (1989)).

[0057] Methods for constructing cDNA libraries in a suitable vector such as a plasmid or phage for propagation in procaryotic or eucaryotic cells are well known to those skilled in the art. [See e.g. Maniatis *et al. Supra*]. Suitable cloning vectors are well known and are widely available.

[0058] The TRAILLK-2 gene or fragment thereof can be isolated a tissue in which said gene is expressed, for example, breast tissue. In one method, mRNA is isolated from a suitable tissue, and first strand cDNA synthesis is carried out. A second round of DNA synthesis can be carried out for the production of the second strand. If desired, the double-stranded cDNA can be cloned into any suitable vector, for example, a plasmid, thereby forming a cDNA library. Oligonucleotide primers targeted to any suitable region of SEQ ID NO:1 can be used for PCR amplification of TRAILLK-2. See e.g. PCR Protocols: A Guide to Method and Application, Ed. M. Innis *et al.*, Academic Press (1990). The PCR amplification comprises template DNA, suitable enzymes, primers, and buffers, and is conveniently carried out in a DNA Thermal Cycler (Perkin Elmer Cetus, Norwalk, CT). A positive result is determined by detecting an appropriately-sized DNA fragment following agarose gel electrophoresis.

Protein Production Methods

[0059] One embodiment of the present invention relates to the substantially purified protein encoded by the TRAILLK-2 gene.

[0060] Skilled artisans will recognize that the proteins of the present invention can be synthesized by a number of different methods, such as chemical methods well known in the art, including solid phase peptide synthesis or recombinant methods. Both methods are described in U.S. Patent 4,617,149, incorporated herein by reference.

[0061] The principles of solid phase chemical synthesis of polypeptides are well known in the art and may be found in general texts in the area. See, e.g., H. Dugas and C. Penney, Bioorganic Chemistry (1981) Springer-Verlag, New York, 54-92. For example, peptides may be synthesized by solid-phase methodology utilizing an Applied Biosystems 430A peptide synthesizer (Applied Biosystems, Foster City, CA) and synthesis cycles supplied by Applied Biosystems.

[0062] The proteins of the present invention can also be produced by recombinant DNA methods using the cloned TRAILLK-2 gene. Recombinant methods are preferred if a high yield is desired. Expression of the cloned gene can be carried out in a variety of suitable host cells, well known to those skilled in the art. For this purpose, the TRAILLK-2 gene is introduced into a host cell by any suitable means, well known to those skilled in the art. While chromosomal integration of the cloned gene is within the scope of the present invention, it is preferred that the gene be cloned into a suitable extra-chromosomally maintained expression vector so that the coding region of the TRAILLK-2 gene is operably-linked to a constitutive or inducible promoter.

[0063] The basic steps in the recombinant production of the TRAILLK-2 protein are:

a) constructing a natural, synthetic or semi-synthetic DNA encoding TRAILLK-2 protein;

b) integrating said DNA into an expression vector in a manner suitable for expressing the TRAILLK-2 protein, either alone or as a fusion protein;

c) transforming or otherwise introducing said vector into an appropriate eucaryotic or prokaryotic host cell forming a recombinant host cell,

d) culturing said recombinant host cell in a manner to express the TRAILLK-2 protein; and

e) recovering and substantially purifying the TRAILLK-2 protein by any suitable means, well known to those skilled in the art.

Expressing Recombinant TRAILLK-2 Protein in Procaryotic and Eucaryotic Host Cells

[0064] Procaryotes may be employed in the production of recombinant TRAILLK-2 protein. For example, the *Escherichia coli* K12 strain 294 (ATCC No. 31446) is particularly useful for the prokaryotic expression of foreign proteins. Other strains of *E. coli*, bacilli such as *Bacillus subtilis*, enterobacteriaceae such as *Salmonella typhimurium* or *Serratia marcescans*, various *Pseudomonas* species and other bacteria, such as *Streptomyces*, may also be employed as host

cells in the cloning and expression of the recombinant proteins of this invention.

[0065] Promoter sequences suitable for driving the expression of genes in procaryotes include b-lactamase [e.g. vector pGX2907, ATCC 39344, contains a replicon and b-lactamase gene], lactose systems [Chang et al., *Nature* (London), 275:615 (1978); Goeddel et al., *Nature* (London), 281:544 (1979)], alkaline phosphatase, and the tryptophan (trp) promoter system [vector pATH1 (ATCC 37695)], which is designed to facilitate expression of an open reading frame as a trpE fusion protein under the control of the trp promoter. Hybrid promoters such as the tac promoter (isolatable from plasmid pDR540, ATCC-37282) are also suitable. Still other bacterial promoters, whose nucleotide sequences are generally known, may be ligated to DNA encoding the protein of the instant invention, using linkers or adapters to supply any required restriction sites. Promoters for use in bacterial systems also will contain a Shine-Dalgarno sequence operably-linked to the DNA encoding the desired polypeptides. These examples are illustrative rather than limiting.

[0066] The proteins of this invention may be synthesized either by direct expression or as a fusion protein comprising the protein of interest as a translational fusion with another protein or peptide which may be removable by enzymatic or chemical cleavage. It is often observed in the production of certain peptides in recombinant systems that expression as a fusion protein prolongs the life span, increases the yield of the desired peptide, or provides a convenient means of purifying the protein. This is particularly relevant when expressing mammalian proteins in procaryotic hosts. A variety of peptidases (e.g. enterokinase and thrombin) which cleave a polypeptide at specific sites or digest the peptides from the amino or carboxy termini (e.g. diaminopeptidase) of the peptide chain are known. Furthermore, particular chemicals (e.g. cyanogen bromide) will cleave a polypeptide chain at specific sites. The skilled artisan will appreciate the modifications necessary to the amino acid sequence (and synthetic or semi-synthetic coding sequence if recombinant means are employed) to incorporate site-specific internal cleavage sites. See e.g., P. Carter, "Site Specific Proteolysis of Fusion Proteins", Chapter 13, in Protein Purification: From Molecular Mechanisms to Large Scale Processes, American Chemical Society, Washington, D.C. (1990).

[0067] In addition to procaryotes, a variety of amphibian expression systems such as frog oocytes, and mammalian cell systems can be used. The choice of a particular host cell depends to some extent on the particular expression vector used. Exemplary mammalian host cells suitable for use in the present invention include HepG-2 (ATCC HB 8065), CV-1 (ATCC CCL 70), LC-MK₂ (ATCC CCL 7.1), 3T3 (ATCC CCL 92), CHO-K1 (ATCC CCL 61), HeLa (ATCC CCL 2), RPMI8226 (ATCC CCL 155), H4IIEC3 (ATCC CCL 1600), C1271 (ATCC CCL 1616), HS-Sultan (ATCC CCL 1484), and BHK-21 (ATCC CCL 10), for example.

[0068] A wide variety of vectors are suitable for transforming mammalian host cells. For example, the pSV2-type vectors comprise segments of the simian virus 40 (SV40) genome required for transcription and polyadenylation. A large number of plasmid pSV2-type vectors have been constructed, such as pSV2-gpt, pSV2-neo, pSV2-dhfr, pSV2-hyg, and pSV2-b-globin, in which the SV40 promoter drives transcription of an inserted gene. These vectors are widely available from sources such as the American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, Maryland, 20852, or the Northern Regional Research Laboratory (NRRL), 1815 N. University Street, Peoria, Illinois, 61604.

[0069] Promoters suitable for expression in mammalian cells include the SV40 late promoter, promoters from eukaryotic genes, such as, for example, the estrogen-inducible chicken ovalbumin gene, the interferon genes, the glucocorticoid-inducible tyrosine aminotransferase gene, the thymidine kinase gene promoter, and the promoters of the major early and late adenovirus genes.

[0070] Plasmid pRSVcat (ATCC 37152) comprises portions of a long terminal repeat of the Rous Sarcoma virus, a virus known to infect chickens and other host cells. This long terminal repeat contains a promoter which is suitable for use in the vectors of this invention. H. Gorman et al., *Proc. Nat. Acad. Sci. (USA)*, 79, 6777 (1982). The plasmid pMSVi (NRRL B-15929) comprises the long terminal repeats of the Murine Sarcoma virus, a virus known to infect mouse and other host cells. The mouse metallothionein promoter has also been well characterized for use in eukaryotic host cells and is suitable for use in the present invention. This promoter is present in the plasmid pdBPV-MMTneo (ATCC 37224) which can serve as the starting material for the construction of other plasmids of the present invention.

[0071] Transfection of mammalian cells with vectors can be performed by a plurality of well known processes including, but not limited to, protoplast fusion, calcium phosphate co-precipitation, electroporation and the like. See, e.g., Maniatis et al., *supra*.

[0072] Some viruses also make appropriate vectors. Examples include the adenoviruses, the adeno-associated viruses, the vaccinia virus, the herpes viruses, the baculoviruses, and the rous sarcoma virus, as described in U.S. Patent 4,775,624, incorporated herein by reference.

[0073] Eucaryotic microorganisms such as yeast and other fungi are also suitable host cells. The yeast *Saccharomyces cerevisiae* is the preferred eucaryotic microorganism. Other yeasts such as *Kluyveromyces lactis* and *Pichia pastoris* are also suitable. For expression in *Saccharomyces*, the plasmid YRp7 (ATCC-40053), for example, may be used. See, e.g., L. Stinchcomb et al., *Nature*, 282, 39 (1979); J. Kingsman et al., *Gene*, 7, 141 (1979); S. Tschemper et al., *Gene*, 10, 157 (1980). Plasmid YRp7 contains the TRP1 gene which provides a selectable marker for use in a

trp1 auxotrophic mutant.

Purification of Recombinantly-Produced TRAILLK-2 Protein

[0074] An expression vector carrying the cloned TRAILLK-2 gene is transformed or transfected into a suitable host cell using standard methods. Cells that contain the vector are propagated under conditions suitable for expression of the recombinant TRAILLK-2 protein. For Example, if the recombinant gene has been placed under the control of an inducible promoter, suitable growth conditions would incorporate the appropriate inducer. The recombinantly-produced protein may be purified from cellular extracts of transformed cells by any suitable means.

[0075] In a preferred process for protein purification, the TRAILLK-2 gene is modified at the 5' end to incorporate several histidine residues at the amino terminus of the TRAILLK-2 protein. This "histidine tag" enables a single-step protein purification method referred to as "immobilized metal ion affinity chromatography" (IMAC), essentially as described in U.S. Patent 4,569,794, which hereby is incorporated by reference. The IMAC method enables rapid isolation of substantially pure recombinant TRAILLK-2 protein starting from a crude extract of cells that express a modified recombinant protein, as described above.

Production of Antibodies

[0076] The proteins of this invention and fragments thereof may be used in the production of antibodies. The term "antibody" as used herein describes antibodies, fragments of antibodies (such as, but not limited to, Fab, Fab', Fab₂', and Fv fragments), and chimeric, humanized, veneered, resurfaced, or CDR-grafted antibodies capable of binding antigens of a similar nature as the parent antibody molecule from which they are derived. The instant invention also encompasses single chain polypeptide binding molecules.

[0077] The production of antibodies, both monoclonal and polyclonal, in animals, especially mice, is well known in the art. See, e.g., C. Milstein, Handbook of Experimental Immunology, (Blackwell Scientific Pub., 1986); J. Goding, Monoclonal Antibodies: Principles and Practice, (Academic Press, 1983). For the production of monoclonal antibodies the basic process begins with injecting a mouse, or other suitable animal, with an immunogen. The mouse is subsequently sacrificed and cells taken from its spleen are fused with myeloma cells, resulting in a hybridoma that reproduces *in vitro*. The population of hybridomas is screened to isolate individual clones, each of which secretes a single antibody species, specific for the immunogen. Each antibody obtained in this way is the clonal product of a single B cell.

[0078] Chimeric antibodies are described in U.S. Patent No. 4,816,567, the entire contents of which is herein incorporated by reference. This reference discloses methods and vectors for the preparation of chimeric antibodies. An alternative approach is provided in U.S. Patent No. 4,816,397, the entire contents of which is herein incorporated by reference. This patent teaches co-expression of the heavy and light chains of an antibody in the same host cell.

[0079] The approach of U.S. Patent 4,816,397 has been further refined in European Patent Publication No. 0 239 400. The teachings of this European patent publication are a preferred format for genetic engineering of monoclonal antibodies. In this technology the complementarity determining regions (CDRs) of a human antibody are replaced with the CDRs of a murine monoclonal antibody, thereby converting the specificity of the human antibody to the specificity of a murine antibody.

[0080] Single chain antibodies and libraries thereof are yet another variety of genetically engineered antibody technology that is well known in the art. (See, e.g. R.E. Bird, *et al.*, *Science* 242:423-426 (1988); PCT Publication Nos. WO 88/01649, WO 90/14430, and WO 91/10737. Single chain antibody technology involves covalently joining the binding regions of heavy and light chains to generate a single polypeptide chain. The binding specificity of the intact antibody molecule is thereby reproduced on a single polypeptide chain.

[0081] The antibodies contemplated by this invention are useful in diagnostics, therapeutics or in diagnostic/therapeutic combinations.

[0082] The proteins of this invention or suitable fragments thereof can be used to generate polyclonal or monoclonal antibodies, and various inter-species hybrids, or humanized antibodies, or antibody fragments, or single-chain antibodies. The techniques for producing antibodies are well known to skilled artisans. (See e.g. A.M. Campbell, Monoclonal Antibody Technology: Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Publishers, Amsterdam (1984); Kohler and Milstein, *Nature* 256, 495-497 (1975); Monoclonal Antibodies: Principles & Applications Ed. J.R. Birch & E.S. Lennox, Wiley-Liss, 1995.

[0083] A protein used as an immunogen may be modified or administered in an adjuvant, by subcutaneous or intraperitoneal injection into, for example, a mouse or a rabbit. For the production of monoclonal antibodies, spleen cells from immunized animals are removed, fused with myeloma cells, such as SP2/0-Ag14 cells, and allowed to become monoclonal antibody producing hybridoma cells in the manner known to the skilled artisan. Hybridomas that secrete a desired antibody molecule can be screened by a variety of well known methods, for example ELISA assay, western blot analysis, or radioimmunoassay (Lutz *et al.* *Exp. Cell Res.* 175, 109-124 (1988); Monoclonal Antibodies: Principles

& Applications Ed. J.R. Birch & E.S. Lennox, Wiley-Liss, 1995).

[0084] For some applications labeled antibodies are desirable. Procedures for labeling antibody molecules are widely known, including for example, the use of radioisotopes, affinity labels, such as biotin or avidin, enzymatic labels, for example horseradish peroxidase, and fluorescent labels, such as FITC or rhodamine (See e.g. Enzyme-Mediated Immunoassay, Ed. T. Ngo, H. Lenhoff, Plenum Press 1985; Principles of Immunology and Immunodiagnosics, R.M. Aloisi, Lea & Febiger, 1988).

[0085] Labeled antibodies are useful for a variety of diagnostic applications. In one embodiment the present invention relates to the use of labeled antibodies to detect the presence of TRAILLK-2. Alternatively, the antibodies could be used in a screen to identify potential modulators of TRAILLK-2. For example, in a competitive displacement assay, the antibody or compound to be tested is labeled by any suitable method. Competitive displacement of an antibody from an antibody-antigen complex by a test compound such that a test compound-antigen complex is formed provides a method for identifying compounds that bind TRAILLK-2.

[0086] Other embodiments of the present invention comprise isolated nucleic acid sequences that encode SEQ ID NO:2, or related nucleic acids that are at least about 75% identical to SEQ ID NO:1, or to their complementary sequence, or nucleic acids that hybridize to SEQ ID NO:1 under low or high stringency conditions. Such sequences may come, for example, from paralogous or orthologous genes.

[0087] The TRAILLK-2 gene (viz. SEQ ID NO:1) and related nucleic acid molecules that encode SEQ ID NO:2, or functional fragments thereof, may be produced by chemical synthetic methods. The synthesis of nucleic acids is well known in the art. See, e.g., E.L. Brown, R. Belagaje, M.J. Ryan, and H.G. Khorana, *Methods in Enzymology*, 68:109-151 (1979). Fragments of the DNA sequence corresponding to the TRAILLK-2 gene could be generated using a conventional DNA synthesizing apparatus, such as the Applied Biosystems Model 380A or 380B DNA synthesizers (Applied Biosystems, Inc., 850 Lincoln Center Drive, Foster City, CA 94404) using phosphoramidite chemistry, thereafter ligating the fragments so as to reconstitute the entire gene. Alternatively, phosphotriester chemistry may be employed to synthesize the nucleic acids of this invention. (See, e.g., M.J. Gait, ed., Oligonucleotide Synthesis, A Practical Approach, (1984)).

[0088] In an alternative methodology, namely PCR, the DNA sequences disclosed and described herein, comprising, for example, a portion or all of SEQ ID NO:1 can be produced from a plurality of starting materials. For example, starting with a cDNA preparation (e.g. cDNA library) derived from a tissue that expresses the TRAILLK-2 gene, suitable oligonucleotide primers complementary to SEQ ID NO:1 or to any sub-region therein, are prepared as described in U.S. Patent No. 4,889,818, hereby incorporated by reference. Other suitable protocols for the PCR are disclosed in PCR Protocols: A Guide to Method and Applications, Ed. Michael A. Innis et al., Academic Press, Inc. (1990). Using PCR, any region of the TRAILLK-2 gene can be targeted for amplification such that full or partial length gene sequences may be produced.

[0089] The ribonucleic acids of the present invention may be prepared using polynucleotide synthetic methods discussed *supra*, or they may be prepared enzymatically, for example, using RNA polymerase to transcribe a TRAILLK-2 DNA template.

[0090] The most preferred systems for preparing the ribonucleic acids of the present invention employ the RNA polymerase from the bacteriophage T7 or the bacteriophage SP6. These RNA polymerases are highly specific, requiring the insertion of bacteriophage-specific sequences at the 5' end of the template to be transcribed. See, Maniatis et al., *supra*.

[0091] This invention also provides nucleic acids, RNA or DNA, that are complementary to SEQ ID NO:1 or SEQ ID NO:4, or fragment thereof.

Nucleic Acid Probes

[0092] The present invention also provides probes and primers useful for a variety of molecular biology techniques including, for example, hybridization screens of genomic, subgenomic, or cDNA libraries, as well as hybridization against nucleic acids derived from cell lines or tissues that originate from drug-resistant tumors. Such hybridization screens are useful as methods to identify homologous and/or functionally related sequences from the same or other organisms. A nucleic acid compound comprising SEQ ID NO:1, or a complementary sequence thereof, or a fragment thereof, which is at least 14 base pairs in length, and which will selectively hybridize to human DNA or mRNA encoding TRAILLK-2 protein or fragment thereof, or a functionally related protein, is provided. Preferably, the 14 or more base pair compound is DNA. See e.g. B. Wallace and G. Miyada, "Oligonucleotide Probes for the Screening of Recombinant DNA Libraries," In *Meth. Enzym.*, 152, 432-442, Academic Press (1987).

[0093] Probes and primers can be prepared by enzymatic or recombinant methods, well known to those skilled in the art (See e.g. Sambrook et al. *supra*). A probe may be a single stranded nucleic acid sequence which is complementary in some particular degree to a nucleic acid sequence sought to be detected ("target sequence"). A probe may be labeled with a detectable moiety such as a radio-isotope, antigen, or chemiluminescent moiety. A description of the

use of nucleic acid hybridization as a procedure for the detection of particular nucleic acid sequences is described in U.S. Patent No. 4,851,330 to Kohne, entitled "Method for Detection, Identification and Quantitation of Non-Viral Organisms."

[0094] Having the DNA sequence of the present invention allows preparation of relatively short DNA (or RNA) sequences having the ability to specifically hybridize to gene sequences disclosed herein. In these aspects, nucleic acid probes of an appropriate length are prepared. The ability of such nucleic acid probes to specifically hybridize to a polynucleotide encoding a TRAILLK-2 gene or related sequence lends particular utility in a variety of embodiments. Most importantly, the probes may be used in a variety of assays for detecting the presence of complementary sequences in a given sample.

[0095] In certain embodiments, it is advantageous to use oligonucleotide primers. The sequence of such primers is designed using a polynucleotide of the present invention for use in detecting, amplifying or mutating a defined segment of a gene or polynucleotide that encodes a TRAILLK-2 polypeptide using PCR technology.

[0096] Preferred nucleic acid sequences employed for hybridization studies, or assays, include probe molecules that are complementary to at least an about 14 to an about 70-nucleotide long stretch of a polynucleotide that encodes a TRAILLK-2 polypeptide, such as the nucleotide base sequences designated as SEQ ID NO:1. A length of at least 14 nucleotides helps to ensure that the fragment is of sufficient length to form a duplex molecule that is both stable and selective. Molecules having complementary sequences over stretches greater than 14 bases in length are generally preferred, though in order to increase stability and selectivity of the hybrid. One will generally prefer to design nucleic acid molecules having gene-complementary stretches of 25 to 40 nucleotides, 55 to 70 nucleotides, or even longer where desired. Such fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means, by application of nucleic acid reproduction technology, such as the PCR TM technology of U.S. Pat. No. 4,603,102, herein incorporated by reference, or by excising selected DNA fragments from recombinant plasmids containing appropriate inserts and suitable restriction enzyme sites.

[0097] The following guidelines are useful for designing probes with desirable characteristics. The extent and specificity of hybridization reactions are affected by a number of factors that determine the sensitivity and specificity of a particular probe, whether perfectly complementary to its target or not. The affect of various experimental parameters and conditions are well known to those skilled in the art.

[0098] First, the stability of the probe:target nucleic acid hybrid should be chosen to be compatible with the assay conditions. This may be accomplished by avoiding long A and T rich sequences, by terminating the hybrids with G:C base pairs and by designing a probe with an appropriate T_m (i.e. melting temperature). The melting profile, including the T_m of a hybrid comprising an oligonucleotide and target sequence, may be determined using a Hybridization Protection Assay. The probe should be chosen so that the length and % GC content result in a T_m about 2°-10° C higher than the temperature at which the final assay will be performed. The base composition of the probe is also a significant factor because G-C base pairs exhibit greater thermal stability as compared to A-T base pairs. Thus, hybridization involving complementary nucleic acids of higher G-C content will be more stable at higher temperatures.

[0099] The ionic strength and incubation temperature under which a probe will be used should also be taken into account. It is known that hybridization will increase as the ionic strength of the reaction mixture increases, and that the thermal stability of molecular hybrids will increase with increasing ionic strength. On the other hand, chemical reagents such as formamide, urea, DMSO and alcohols, which disrupt hydrogen bonds, increase the stringency of hybridization. Destabilization of hydrogen bonds by such reagents can greatly reduce the T_m. In general, optimal hybridization for synthetic oligonucleotide probes of about 10-50 bases in length occurs approximately 5° C below the melting temperature for a given duplex. Incubation at temperatures below the optimum may allow mismatched base sequences to hybridize and can therefore result in reduced specificity.

[0100] The length of the target nucleic acid sequence and, accordingly, the length of the probe sequence can also be important. In some cases, there may be several sequences from a particular region, varying in location and length, which will yield probes with the desired hybridization characteristics. In other cases, one sequence may be significantly better than another even though the one sequence differs merely by a single base. Finally, there can be intramolecular and intermolecular hybrids formed within a probe if there is sufficient self-complementarity. Such structures can be avoided through careful probe design. Computer programs are available to search for this type of interaction.

[0101] A probe molecule may be used for hybridizing to a sample suspected of possessing a TRAILLK-2 or TRAILLK-2-related nucleotide sequence. The hybridization reaction is carried out under suitable conditions of stringency.

[0102] Alternatively, such DNA molecules may be used in a number of techniques including their use as: (1) diagnostic tools to detect polymorphisms in DNA samples from a human or other mammal; (2) means for detecting and isolating homologs of TRAILLK-2 and related polypeptides from a DNA library potentially containing such sequences; (3) primers for hybridizing to related sequences for the purpose of amplifying those sequences; and (4) primers for altering the native TRAILLK-2 DNA sequences; as well as other techniques which rely on the similarity of the DNA sequences to those of the TRAILLK-2 DNA segments herein disclosed.

[0103] Once synthesized, oligonucleotide probes may be labeled by any of several well known methods. See e.g.

Maniatis *et al.*, Molecular Cloning (2d ed. 1989). Useful labels include radioisotopes, as well as non-radioactive reporting groups. Isotopic labels include H³, S³⁵, P³², I¹²⁵, Cobalt, and C¹⁴. Most methods of isotopic labeling involve the use of enzymes and include methods such as nick-translation, end-labeling, second strand synthesis, and reverse transcription. When using radio-labeled probes, hybridization can be detected by autoradiography, scintillation counting, or gamma counting. The detection method selected will depend upon the hybridization conditions and the particular radio isotope used for labeling.

[0104] Non-isotopic materials can also be used for labeling, and may be introduced internally into the sequence or at the end of the sequence. Modified nucleotides may be incorporated enzymatically or chemically, and chemical modifications of the probe may be performed during or after synthesis of the probe, for example, by the use of non-nucleotide linker groups. Non-isotopic labels include fluorescent molecules, chemiluminescent molecules, enzymes, cofactors, enzyme substrates, haptens or other ligands. In a preferred embodiment of the invention, the length of an oligonucleotide probe is greater than or equal to about 18 nucleotides and less than or equal to about 50 nucleotides. Labeling of an oligonucleotide of the present invention may be performed enzymatically using [³²P]-labeled ATP and the enzyme T4 polynucleotide kinase.

Vectors

[0105] Another aspect of the present invention relates to recombinant DNA cloning vectors and expression vectors comprising the nucleic acids of the present invention. The preferred nucleic acid vectors are those which comprise DNA. The most preferred recombinant DNA vectors comprise the isolated DNA sequence, SEQ ID NO:1.

[0106] The skilled artisan understands that choosing the most appropriate cloning vector or expression vector depends upon a number of factors including the availability of restriction enzyme sites, the type of host cell into which the vector is to be transfected or transformed, the purpose of the transfection or transformation (e.g., stable transformation as an extrachromosomal element, or integration into the host chromosome), the presence or absence of readily assayable or selectable markers (e.g., antibiotic resistance and metabolic markers of one type and another), and the number of copies of the gene desired in the host cell.

[0107] Vectors suitable to carry the nucleic acids of the present invention comprise RNA viruses, DNA viruses, lytic bacteriophages, lysogenic bacteriophages, stable bacteriophages, plasmids, viroids, and the like. The most preferred vectors are plasmids.

[0108] When preparing an expression vector the skilled artisan understands that there are many variables to be considered, for example, whether to use a constitutive or inducible promoter. The practitioner also understands that the amount of nucleic acid or protein to be produced dictates, in part, the selection of the expression system. Regarding promoter sequences, inducible promoters are preferred because they enable high level, regulatable expression of an operably-linked gene. The skilled artisan will recognize a number of suitable promoters that respond to a variety of inducers, for example, carbon source, metal ions, and heat. Other relevant considerations regarding an expression vector include whether to include sequences for directing the localization of a recombinant protein. For example, a sequence encoding a signal peptide preceding the coding region of a gene is useful for directing the extracellular export of a resulting polypeptide.

[0109] The present invention also provides a method for constructing a recombinant host cell capable of expressing proteins comprising SEQ ID NO:2, said method comprising transforming or otherwise introducing into a host cell a recombinant DNA vector that comprises an isolated DNA sequence that encodes SEQ ID NO:2. A suitable host cell is any eucaryotic cell that can accommodate high level expression of an exogenously introduced gene or protein, and that will incorporate said protein into its membrane structure. Vectors for expression are those which comprise SEQ ID NO:1 or fragment thereof. Transformed host cells may be cultured under conditions well known to skilled artisans such that SEQ ID NO:2 is expressed, thereby producing a recombinant TRAILLK-2 protein in the recombinant host cell.

[0110] For the purpose of identifying compounds having utility as modifiers of apoptosis, it would be desirable to identify compounds that bind the TRAILLK-2 protein and/or modify its activity. For the purpose of identifying or developing inhibitors or other modifiers that, for example, activate or inhibit the proteins disclosed herein, it would be desirable to identify compounds that bind the TRAILLK-2 protein. A method for determining agents that bind the TRAILLK-2 protein comprises contacting the TRAILLK-2 protein with a test compound and monitoring binding by any suitable means.

[0111] The instant invention provides a screening system for discovering compounds that bind the TRAILLK-2 protein, said screening system comprising the steps of:

- a) preparing TRAILLK-2 protein;
- b) exposing said TRAILLK-2 protein to a test compound;

c) quantifying the binding of said compound to TRAILLK-2 protein by any suitable means.

[0112] Utilization of the screening system described above provides a means to determine compounds that may alter the biological function of TRAILLK-2. This screening method may be adapted to large-scale, automated procedures such as a PANDEX® (Baxter-Dade Diagnostics) system, allowing for efficient high-volume screening of potential therapeutic agents.

[0113] In such a screening protocol TRAILLK-2 is prepared as described herein, preferably using recombinant DNA technology. A test compound is introduced into a reaction vessel containing the TRAILLK-2 protein or fragment thereof. Binding of TRAILLK-2 by a test compound is determined by any suitable means. For example, in one method radioactively-labeled or chemically-labeled test compound may be used. Binding of the protein by the compound is assessed, for example, by quantifying bound label versus unbound label using any suitable method. Binding of a test compound may also be carried out by a method disclosed in U.S. Patent 5,585,277, which hereby is incorporated by reference. In this method, binding of a test compound to a protein is assessed by monitoring the ratio of folded protein to unfolded protein, for example by monitoring sensitivity of said protein to a protease, or amenability to binding of said protein by a specific antibody against the folded state of the protein.

[0114] The foregoing screening methods are useful for identifying a ligand of a TRAILLK-2 protein, as a lead to a pharmaceutical compound for modulating apoptosis. A ligand that binds TRAILLK-2, or related fragment thereof, is identified, for example, by combining a test ligand with TRAILLK-2 under conditions that cause the protein to exist in a ratio of folded to unfolded states. If the test ligand binds the folded state of the protein, the relative amount of folded protein will be higher than in the case of a test ligand that does not bind the protein. The ratio of protein in the folded versus unfolded state is easily determinable by, for example, susceptibility to digestion by a protease, or binding to a specific antibody, or binding to chaperonin protein, or binding to any suitable surface.

[0115] The present invention also provides a pharmaceutical composition comprising as the active agent a polypeptide compound represented by SEQ ID NO:2, or a pharmaceutically acceptable non-toxic salt thereof, and a pharmaceutically acceptable solid or liquid carrier. For example, compounds comprising TRAILLK-2 can be admixed with conventional pharmaceutical carriers and excipients, and used in the form of tablets, capsules, elixirs, suspensions, syrups, waters, and the like. The compositions comprising TRAILLK-2 will contain from about 0.1% to 90% by weight of the active compound, and more generally from about 10% to 30%. The compositions may contain common carriers and excipients such as corn starch or gelatin, lactose, sucrose, microcrystalline cellulose, kaolin, mannitol, dicalcium phosphate, sodium chloride, and alginate acid. The compounds can be formulated for oral or parenteral administration.

[0116] Skilled artisans will recognize that IC_{50} values are dependent on the selectivity of the compound tested. For example, a compound with an IC_{50} which is less than 10 nM is generally considered an excellent candidate for drug therapy. However, a compound which has a lower affinity, but is selective for a particular target, may be an even better candidate. The skilled artisan will recognize that any information regarding the binding potential, inhibitory activity, or selectivity of a particular compound is useful toward the development of pharmaceutical products.

[0117] The following examples more fully describe the present invention. Those skilled in the art will recognize that the particular reagents, equipment, and procedures described are merely illustrative and are not intended to limit the present invention in any manner.

EXAMPLE 1

RT-PCR Amplification of TRAILLK-2 Gene from mRNA

[0118] A TRAILLK-2 gene is isolated by reverse transcriptase PCR (RT-PCR) using conventional methods. Total RNA from a tissue that expresses the TRAILLK-2 gene, is prepared using standard methods. First strand cDNA synthesis is achieved using a commercially available kit (SuperScript™ System; Life Technologies) by PCR in conjunction with specific primers directed at any suitable region of SEQ ID NO:1, for example, at the ATG start site at base pair 49, and at the TGA stop site after base pair 798.

[0119] Amplification is carried out by adding to the first strand cDNA (dried under vacuum): 8 μ l of 10X synthesis buffer (200 mM Tris-HCl, pH 8.4; 500 mM KCl, 25 mM $MgCl_2$, 1 μ g/ μ l BSA); 68 μ l distilled water; 1 μ l each of a 10 μ M solution of each primer; and 1 μ l Taq DNA polymerase (2 to 5 U/ μ l). The reaction is heated at 94° C for 5 min. to denature the RNA/cDNA hybrid. Then, 15 to 30 cycles of PCR amplification are performed using any suitable thermal cycle apparatus. The amplified sample may be analyzed by agarose gel electrophoresis to check for an appropriately-sized fragment.

EXAMPLE 2

Production of a Vector for Expressing TRAILLK-2 in a Host Cell

[0120] An expression vector suitable for expressing TRAILLK-2 or fragment thereof in a variety of procaryotic host cells, such as *E. coli* is easily made. The vector contains an origin of replication (Ori), an ampicillin resistance gene (Amp) useful for selecting cells which have incorporated the vector following a transformation procedure, and further comprises the T7 promoter and T7 terminator sequences in operable linkage to a TRAILLK-2 coding region. Plasmid pET11A (obtained from Novogen, Madison WI) is a suitable parent plasmid. pET11A is linearized by restriction with endonucleases NdeI and BamHI. Linearized pET11A is ligated to a DNA fragment bearing NdeI and BamHI sticky ends and comprising the coding region of the TRAILLK-2 gene as disclosed by SEQ ID NO:1 or fragment thereof.

[0121] The TRAILLK-2 gene used in this construction may be slightly modified at the 5' end (amino terminus of encoded protein) in order to simplify purification of the encoded protein product. For this purpose, an oligonucleotide encoding 8 histidine residues is inserted after the ATG start codon. Placement of the histidine residues at the amino terminus of the encoded protein serves to enable the IMAC one-step protein purification procedure.

EXAMPLE 3

Recombinant Expression and Purification of TRAILLK-2 Protein

[0122] An expression vector that carries an ORF encoding TRAILLK-2 or fragment thereof and which ORF is operably-linked to an expression promoter is transformed into *E. coli* BL21 (DE3) (*hsdS gal lclts857 ind1 Sam7 nin5 lacUV5-T7 gene 1*) using standard methods. Transformants, selected for resistance to ampicillin, are chosen at random and tested for the presence of the vector by agarose gel electrophoresis using quick plasmid preparations. Colonies which contain the vector are grown in L broth and the protein product encoded by the vector-borne ORF is purified by immobilized metal ion affinity chromatography (IMAC), essentially as described in US Patent 4,569,794.

[0123] Briefly, the IMAC column is prepared as follows. A metal-free chelating resin (e.g. Sepharose 6B IDA, Pharmacia) is washed in distilled water to remove preservative substances and infused with a suitable metal ion [e.g. Ni(II), Co(II), or Cu(II)] by adding a 50mM metal chloride or metal sulfate aqueous solution until about 75% of the interstitial spaces of the resin are saturated with colored metal ion. The column is then ready to receive a crude cellular extract containing the recombinant protein product.

[0124] After removing unbound proteins and other materials by washing the column with any suitable buffer, pH 7.5, the bound protein is eluted in any suitable buffer at pH 4.3, or preferably with an imidazole-containing buffer at pH 7.5.

EXAMPLE 4

Tissue Distribution of TRAILLK-2 mRNA

[0125] The presence of TRAILLK-2 mRNA in a variety of human tissues was analyzed by Northern analysis. Total RNA from different tissues or cultured cells was isolated by a standard guanidine chloride/phenol extraction method, and poly-A⁺ RNA was isolated using oligo(dT)-cellulose type 7 (Pharmacia). Electrophoresis of RNA samples was carried out in formaldehyde followed by capillary transfer to Zeta-Probe TM nylon membranes (Bio-Rad, Hercules, Calif.). SEQ ID NO:1 was the template for generating probes using a MultiPrime™ random priming kit (Amersham, Arlington Heights, Ill.). The efficiency of the labeling reaction was approximately 4×10^{10} cpm incorporated per μ g of template. The hybridization buffer contained 0.5M sodium phosphate, 7% SDS (wt/vol), 1% BSA (wt/vol), and 1 mM EDTA. Prehybridization was carried out in hybridization buffer at 65° C for 2 h and ³²P-labeled probe was added and incubation continued overnight. The filters were washed in Buffer A (40 mM sodium phosphate pH 7.2, 5% SDS [wt/vol], 0.5% BSA [wt/vol], and 1 mM EDTA) at 65° C for 1 h, and then in Buffer B (40 mM sodium phosphate, pH 7.2, 1% SDS [wt/vol], and 1 mM EDTA) at 65° C for 20 min. The filters were air-dried and exposed to Kodak X-OMAT AR film at -80° C with an intensifying screen.

EXAMPLE 5

Detecting Ligands that Bind TRAILLK-2 Using a Chaperonin Protein Assay

[0126] The wells of an ELISA plate are coated with chaperonin by incubation for several hours with a 4 μ g/ml solution of the protein in Tris-buffered Saline (TBS: 10 mM Tris-HCl, pH7.5, 0.2M NaCl). The plates are then washed 3 times with TBS containing 0.1% Tween-20 (TBST). Then, a mixture of TRAILLK-2 protein (sufficient amount to saturate about

50% of the binding sites on chaperonin) and test compound (10^{-9} to 10^{-5} M) in about 50 μ l volume is added to each well of the plate for an incubation of about 60 minutes. Aliquots of the well solutions are then transferred to the wells of fresh plates and incubated for 60 minutes at room temperature, followed by 3 washes with TBST. Next, about 50 μ l of an antibody specific for TRAILLK-2 plus 5% nonfat dry milk are added to each well for a 30 minute incubation at room temperature. After washing, about 50 μ l of goat anti-rabbit IgG alkaline phosphatase conjugate at an appropriate dilution in TBST plus 5% nonfat dry milk are added to each well and incubated 30 minutes at room temperature. The plates are washed again with TBST and 0.1 ml of 1 mg/ml p-nitrophenylphosphate in 0.1% diethanolamine is added. Color development (proportional to bound alkaline phosphatase antibody conjugate) is monitored with an ELISA plate reader. When test ligand binding has occurred, ELISA analysis reveals TRAILLK-2 in solution at higher concentrations than in the absence of test ligand.

EXAMPLE 6

Production of an Antibody to TRAILLK-2 Protein

[0127] Substantially pure TRAILLK-2 protein or fragment thereof is isolated from transfected or transformed cells using any of the well known methods in the art, or by a method specifically disclosed herein. Concentration of protein in a final preparation is adjusted, for example, by filtration through an Amicon filter device such that the level is about 1 to 5 μ g/ml. Monoclonal or polyclonal antibody can be prepared as follows.

[0128] Monoclonal antibody can be prepared from murine hybridomas according to the method of Kohler and Milstein (*Nature*, 256, 495, 1975), or a modified method thereof. Briefly, a mouse is repetitively inoculated with a few micrograms of the protein or fragment thereof, or fusion peptide thereof, over a period of a few weeks. The mouse is then sacrificed and the antibody producing cells of the spleen isolated. The spleen cells are fused by means of polyethylene glycol with mouse myeloma cells. Fused cells that produce antibody are identified by any suitable immunoassay, for example, ELISA, as described in E. Engvall, *Meth. Enzymol.*, 70, 419, 1980.

[0129] Polyclonal antiserum can be prepared by well known methods (See e. g. J. Vaitukaitis *et al. Clin. Endocrinol. Metab.* 33, 988, 1971) that involve immunizing suitable animals with the proteins, fragments thereof, or fusion proteins thereof, disclosed herein. Small doses (e.g. nanogram amounts) of antigen administered at multiple intradermal sites appears to be the most reliable method

Annex to the description

[0130]

SEQUENCE LISTING

SEQ ID NO:1

GAAAGGGAGCAGTCACGCCTTACTTCTTGCCTTAAGAAAAGAGAAGAAATGAAACTGAA
GGAGTGTGTTTCCATCCTCCACGGAAGGAAAGCCCCTCTGTCCGATCCTCCAAAGACG
GAAAGCTCCCACGCGTCCGGCTGGCTGCAACCTTGCTGCTGGCACTGCTGTCTTGCTG
CCTCAANGGTGGTGTCTTTCTACCAGGTGGCCGCCCTGCAAGGGGACCTGGCCAGCCT
CCGGGCAGAGCTGCAGGGCCACCACGCGGAGAAGCTGCCAGCAGGAGCAGGAGCCCC
CCAAGGCCGGCCTGGAGGAAGCTCCCAGCTGTCACCGCGGACTGAAAATCTTTGAAC
CACCAGCTCCAGGAGAAGGCAACTCCAGTCAGAACAGCAGAAATAAGCGTGCCGTTCA
GGGTCCAGAAGAAAACAGGATCTTACACATTTGTTCCATGGCTTCTCAGCTTTAAAAGGGG
AAGTGCCCTAGAAGAAAAAGAGAATAAAATATTGGTCAAAGAAACTGGTTACTTTTTATA
TATGGTCAGGTTTTATATACTGATAAGACCTACGCCATGGGACATCTAATTCAGAGGAAG
AAGGTCCATGTCTTTGGGGATGAATTGAGTCTGGTGACTTTGTTTCGATGTATTCAAAT
ATGCCTGAAACACTACCCAATAATTCCTGCTATTGAGCTGGCATTGCAAACTGGAAGAA
GGAGATGAACTCCAATTGCAATACCAAGAGAAAAATGCACAAATATCACTGGATGGAGAT
GTCACATTTTTTGGTGCATTGAAACTGCTGTGACCTACTTACACCATGTCTGTAGCTATTT
TCCTCCCTTTCTCTGTACCTCTAAGAAGAAAGAATCTAACTG

SEQ ID NO:2

MKLKECVSILPRKESPSVRSSKDGKLPRVRLAATLLALLSCCLXGGVFLPGGRPARGPGQP
PGRAAGPPRGEAASRSRSPRPAPWRKLPVAVTAGLKIFEPAPGEGNSSQNSRNKRAVQGP
EETGSYTFVPWLLSFKRGSAL EKENKILVKETGYFFIYGQVLYTDKTYAMGHLIQRKKVHVF
GDELSLVTLFRCIQNMPETLPNNSCYSAGIAKLEEGDELQLAIPRENAQISLDGDVTFFGALKL
L

SEQUENCE LISTING

5 <110> Song, Ho Yeong

10 <120> TNF Ligand Family Gene

<130> X11834

15 <140>

<141>

20 <160> 2

25 <170> PatentIn Ver. 2.0

<210> 1

30 <211> 871

<212> DNA

35 <213> Homo sapiens

<220>

40 <221> CDS

<222> (49)..(798)

45 <400> 1

gaaagggagc agtcacgcct tactttcttgc cttaagaaaa gagaagaa atg aaa ctg 57

Met Lys Leu

50 1

aag gag tgt gtt tcc atc ctc cca cgg aag gaa agc ccc tct gtc cga 105

Lys Glu Cys Val Ser Ile Leu Pro Arg Lys Glu Ser Pro Ser Val Arg

5 5 10 15

10 20 25 30 35

15 40 45 50

20 55 60 65

25 70 75 80

30 85 90 95

35 100 105 110 115

40 120 125 130

45 135 140 145 150

50 155 160 165 170

55 175 180 185 190

195 200 205 210 215

220 225 230 235 240

245 250 255 260 265

270 275 280 285 290

295 300 305 310 315

320 325 330 335 340

345 350 355 360 365

370 375 380 385 390

395 400 405 410 415

420 425 430 435 440

445 450 455 460 465

470 475 480 485 490

495 500 505 510 515

520 525 530 535 540

545 550 555 560 565

570 575 580 585 590

595 600 605 610 615

620 625 630 635 640

645 650 655 660 665

670 675 680 685 690

695 700 705 710 715

720 725 730 735 740

745 750 755 760 765

770 775 780 785 790

795 800 805 810 815

820 825 830 835 840

845 850 855 860 865

870 875 880 885 890

895 900 905 910 915

920 925 930 935 940

945 950 955 960 965

970 975 980 985 990

995 1000 1005 1010 1015

1020 1025 1030 1035 1040

1045 1050 1055 1060 1065

1070 1075 1080 1085 1090

1095 1100 1105 1110 1115

1120 1125 1130 1135 1140

1145 1150 1155 1160 1165

1170 1175 1180 1185 1190

1195 1200 1205 1210 1215

1220 1225 1230 1235 1240

1245 1250 1255 1260 1265

1270 1275 1280 1285 1290

1295 1300 1305 1310 1315

1320 1325 1330 1335 1340

1345 1350 1355 1360 1365

1370 1375 1380 1385 1390

1395 1400 1405 1410 1415

1420 1425 1430 1435 1440

1445 1450 1455 1460 1465

1470 1475 1480 1485 1490

1495 1500 1505 1510 1515

1520 1525 1530 1535 1540

1545 1550 1555 1560 1565

1570 1575 1580 1585 1590

1595 1600 1605 1610 1615

1620 1625 1630 1635 1640

1645 1650 1655 1660 1665

1670 1675 1680 1685 1690

1695 1700 1705 1710 1715

1720 1725 1730 1735 1740

1745 1750 1755 1760 1765

1770 1775 1780 1785 1790

1795 1800 1805 1810 1815

1820 1825 1830 1835 1840

1845 1850 1855 1860 1865

1870 1875 1880 1885 1890

1895 1900 1905 1910 1915

1920 1925 1930 1935 1940

1945 1950 1955 1960 1965

1970 1975 1980 1985 1990

1995 2000 2005 2010 2015

2020 2025 2030 2035 2040

2045 2050 2055 2060 2065

2070 2075 2080 2085 2090

2095 2100 2105 2110 2115

2120 2125 2130 2135 2140

2145 2150 2155 2160 2165

2170 2175 2180 2185 2190

2195 2200 2205 2210 2215

2220 2225 2230 2235 2240

2245 2250 2255 2260 2265

2270 2275 2280 2285 2290

2295 2300 2305 2310 2315

2320 2325 2330 2335 2340

2345 2350 2355 2360 2365

2370 2375 2380 2385 2390

2395 2400 2405 2410 2415

2420 2425 2430 2435 2440

2445 2450 2455 2460 2465

2470 2475 2480 2485 2490

2495 2500 2505 2510 2515

2520 2525 2530 2535 2540

2545 2550 2555 2560 2565

2570 2575 2580 2585 2590

2595 2600 2605 2610 2615

2620 2625 2630 2635 2640

2645 2650 2655 2660 2665

2670 2675 2680 2685 2690

2695 2700 2705 2710 2715

2720 2725 2730 2735 2740

2745 2750 2755 2760 2765

2770 2775 2780 2785 2790

2795 2800 2805 2810 2815

2820 2825 2830 2835 2840

2845 2850 2855 2860 2865

2870 2875 2880 2885 2890

2895 2900 2905 2910 2915

2920 2925 2930 2935 2940

2945 2950 2955 2960 2965

2970 2975 2980 2985 2990

2995 3000 3005 3010 3015

3020 3025 3030 3035 3040

3045 3050 3055 3060 3065

3070 3075 3080 3085 3090

3095 3100 3105 3110 3115

3120 3125 3130 3135 3140

3145 3150 3155 3160 3165

3170 3175 3180 3185 3190

3195 3200 3205 3210 3215

3220 3225 3230 3235 3240

3245 3250 3255 3260 3265

3270 3275 3280 3285 3290

3295 3300 3305 3310 3315

3320 3325 3330 3335 3340

3345 3350 3355 3360 3365

3370 3375 3380 3385 3390

3395 3400 3405 3410 3415

3420 3425 3430 3435 3440

3445 3450 3455 3460 3465

3470 3475 3480 3485 3490

3495 3500 3505 3510 3515

3520 3525 3530 3535 3540

3545 3550 3555 3560 3565

3570 3575 3580 3585 3590

3595 3600 3605 3610 3615

3620 3625 3630 3635 3640

3645 3650 3655 3660 3665

3670 3675 3680 3685 3690

3695 3700 3705 3710 3715

3720 3725 3730 3735 3740

3745 3750 3755 3760 3765

3770 3775 3780 3785 3790

3795 3800 3805 3810 3815

3820 3825 3830 3835 3840

3845 3850 3855 3860 3865

3870 3875 3880 3885 3890

3895 3900 3905 3910 3915

3920 3925 3930 3935 3940

3945 3950 3955 3960 3965

3970 3975 3980 3985 3990

3995 4000 4005 4010 4015

4020 4025 4030 4035 4040

4045 4050 4055 4060 4065

4070 4075 4080 4085 4090

4095 4100 4105 4110 4115

4120 4125 4130 4135 4140

4145 4150 4155 4160 4165

4170 4175 4180 4185 4190

4195 4200 4205 4210 4215

4220 4225 4230 4235 4240

4245 4250 4255 4260 4265

4270 4275 4280 4285 4290

4295 4300 4305 4310 4315

4320 4325 4330 4335 4340

4345 4350 4355 4360 4365

4370 4375 4380 4385 4390

4395 4400 4405 4410 4415

4420 4425 4430 4435 4440

4445 4450 4455 4460 4465

4470 4475 4480 4485 4490

4495 4500 4505 4510 4515

4520 4525 4530 4535 4540

4545 4550 4555 4560 4565

4570 4575 4580 4585 4590

4595 4600 4605 4610 4615

4620 4625 4630 4635 4640

4645 4650 4655 4660 4665

4670 4675 4680 4685 4690

4695 4700 4705 4710 4715

4720 4725 4730 4735 4740

4745 4750 4755 4760 4765

4770 4775 4780 4785 4790

4795 4800 4805 4810 4815

4820 4825 4830 4835 4840

4845 4850 4855 4860 4865

4870 4875 4880 4885 4890

4895 4900 4905 4910 4915

4920 4925 4930 4935 4940

4945 4950 4955 4960 4965

4970 4975 4980 4985 4990

4995 5000 5005 5010 5015

5020 5025 5030 5035 5040

5045 5050 5055 5060 5065

5070 5075 5080 5085 5090

5095 5100 5105 5110 5115

5120 5125 5130 5135 5140

5145 5150 5155 5160 5165

5170 5175 5180 5185 5190

5195 5200 5205 5210 5215

5220 5225 5230 5235 5240

5245 5250 5255 5260 5265

5270 5275 5280 5285 5290

5295 5300 5305 5310 5315

5320 5325 5330 5335 5340

5345 5350 5355 5360 5365

5370 5375 5380 5385 5390

5395 5400 5405 5410 5415

5420 5425 5430 5435 5440

5445 5450 5455 5460 5465

5470 5475 5480 5485 5490

5495 5500 5505 5510 5515

5520 5525 5530 5535 5540

5545 5550 5555 5560 5565

5570 5575 5580 5585 5590

5595 5600 5605 5610 5615

5620 5625 5630 5635 5640

5645 5650 5655 5660 5665

5670 5675 5680 5685 5690

5695 5700 5705 5710 5715

5720 5725 5730 5735 5740

5745 5750 5755 5760 5765

5770 5775 5780 5785 5790

5795 5800 5805 5810 5815

5820 5825 5830 5835 5840

5845 5850 5855 5860 5865

5870 5875 5880 5885 5890

5895 5900 5905 5910 5915

5920 5925 5930 5935 5940

5945 5950 5955 5960 5965

5970 5975 5980 5985 5990

5995 6000 6005 6010 6015

6020 6025 6030 6035 6040

6045 6050 6055 6060 6065

6070 6075 6080 6085 6090

6095 6100 6105 6110 6115

6120 6125 6130 6135 6140

6145 6150 6155 6160 6165

6170 6175 6180 6185 6190

6195 6200 6205 6210 6215

6220 6225 6230 6235 6240

6245 6250 6255 6260 6265

6270 6275 6280 6285 6290

6295 6300 6305 6310 6315

6320 6325 6330 6335 6340

6345 6350 6355 6360 6365

6370 6375 6380 6385 6390

6395 6400 6405 6410 6415

6420 6425 6430 6435 6440

6445 6450 6455 6460 6465

6470 6475 6480 6485 6490

6495 6500 6505 6510 6515

6520 6525 6530 6535 6540

6545 6550 6555 6560 6565

6570 6575 6580 6585 6590

6595 6600 6605 6610 6615

6620 6625 6630 6635 6640

6645 6650 6655 6660 6665

6670 6675 6680 6685 6690

6695 6700 6705 6710 6715

6720 6725 6730 6735 674

cca tgg ctt ctc agc ttt aaa agg gga agt gcc cta gaa gaa aaa gag 489
 Pro Trp Leu Leu Ser Phe Lys Arg Gly Ser Ala Leu Glu Glu Lys Glu
 5 135 140 145

aat aaa ata ttg gtc aaa gaa act ggt tac ttt ttt ata tat ggt cag 537
 Asn Lys Ile Leu Val Lys Glu Thr Gly Tyr Phe Phe Ile Tyr Gly Gln
 10 150 155 160

gtt tta tat act gat aag acc tac gcc atg gga cat cta att cag agg 585
 Val Leu Tyr Thr Asp Lys Thr Tyr Ala Met Gly His Leu Ile Gln Arg
 15 165 170 175

aag aag gtc cat gtc ttt ggg gat gaa ttg agt ctg gtg act ttg ttt 633
 Lys Lys Val His Val Phe Gly Asp Glu Leu Ser Leu Val Thr Leu Phe
 20 180 185 190 195

cga tgt att caa aat atg cct gaa aca cta ccc aat aat tcc tgc tat 681
 Arg Cys Ile Gln Asn Met Pro Glu Thr Leu Pro Asn Asn Ser Cys Tyr
 25 200 205 210

tca gct ggc att gca aaa ctg gaa gaa gga gat gaa ctc caa ctt gca 729
 Ser Ala Gly Ile Ala Lys Leu Glu Glu Gly Asp Glu Leu Gln Leu Ala
 30 215 220 225

ata cca aga gaa aat gca caa ata tca ctg gat gga gat gtc aca ttt 777
 Ile Pro Arg Glu Asn Ala Gln Ile Ser Leu Asp Gly Asp Val Thr Phe
 35 230 235 240

ttt ggt gca ttg aaa ctg ctg tgacctactt acaccatgtc tgtagctatt 828

Phe Gly Ala Leu Lys Leu Leu

245

250

ttctccctt tctctgtacc tctaagaaga aagaatctaa ctg

871

<210> 2

<211> 250

<212> PRT

<213> Homo sapiens

<400> 2

Met Lys Leu Lys Glu Cys Val Ser Ile Leu Pro Arg Lys Glu Ser Pro

1

5

10

15

Ser Val Arg Ser Ser Lys Asp Gly Lys Leu Pro Arg Val Arg Leu Ala

20

25

30

Ala Thr Leu Leu Leu Ala Leu Leu Ser Cys Cys Leu Xaa Gly Gly Val

35

40

45

Phe Leu Pro Gly Gly Arg Pro Ala Arg Gly Pro Gly Gln Pro Pro Gly

50

55

60

Arg Ala Ala Gly Pro Pro Arg Gly Glu Ala Ala Ser Arg Ser Arg Ser

65

70

75

80

Pro Pro Arg Pro Ala Trp Arg Lys Leu Pro Ala Val Thr Ala Gly Leu

85

90

95

EP 0 921 194 A2

	Lys	Ile	Phe	Glu	Pro	Pro	Ala	Pro	Gly	Glu	Gly	Asn	Ser	Ser	Gln	Asn
					100				105						110	
5	"															
	Ser	Arg	Asn	Lys	Arg	Ala	Val	Gln	Gly	Pro	Glu	Glu	Thr	Gly	Ser	Tyr
					115				120					125		
10	"															
	Thr	Phe	Val	Pro	Trp	Leu	Leu	Ser	Phe	Lys	Arg	Gly	Ser	Ala	Leu	Glu
15	"				130				135					140		
	"															
	Glu	Lys	Glu	Asn	Lys	Ile	Leu	Val	Lys	Glu	Thr	Gly	Tyr	Phe	Phe	Ile
20	"				145				150				155			160
	"															
	Tyr	Gly	Gln	Val	Leu	Tyr	Thr	Asp	Lys	Thr	Tyr	Ala	Met	Gly	His	Leu
25	"								165				170			175
	"															
	Ile	Gln	Arg	Lys	Lys	Val	His	Val	Phe	Gly	Asp	Glu	Leu	Ser	Leu	Val
30	"								180				185			190
	"															
	Thr	Leu	Phe	Arg	Cys	Ile	Gln	Asn	Met	Pro	Glu	Thr	Leu	Pro	Asn	Asn
35	"															
	"				195				200						205	
	"															
40	Ser	Cys	Tyr	Ser	Ala	Gly	Ile	Ala	Lys	Leu	Glu	Glu	Gly	Asp	Glu	Leu
	"								210				215			220
	"															
45	Gln	Leu	Ala	Ile	Pro	Arg	Glu	Asn	Ala	Gln	Ile	Ser	Leu	Asp	Gly	Asp
	"															
	"								225				230			235
	"															240
50	"															
	Val	Thr	Phe	Phe	Gly	Ala	Leu	Lys	Leu	Leu						
	"															
	"								245						250	
	"															
55	"															

Claims

1. A substantially pure protein having the amino acid sequence which is SEQ ID NO: 2.
- 5 2. An isolated nucleic acid compound encoding the protein of Claim 1, said protein having the amino acid sequence that is SEQ ID NO: 2.
3. An isolated nucleic acid compound that encodes a protein having TNF family ligand activity wherein said nucleic acid hybridizes to SEQ ID NO:1 under high stringency conditions.
- 10 4. A vector comprising an isolated nucleic acid compound of Claim 2 or Claim 3.
5. A vector, as in Claim 4, wherein said isolated nucleic acid compound is operably-linked to a promoter sequence.
- 15 6. A host cell containing a vector of Claim 4 or Claim 5.
7. A method for constructing a recombinant host cell having the potential to express SEQ ID NO:2, said method comprising introducing into said host cell by any suitable means a vector of Claim 5.
- 20 8. A method for identifying compounds that bind a protein identified herein as SEQ ID NO:2, comprising the steps of:
 - a) admixing a substantially purified preparation of a protein comprising SEQ ID NO:2 with a test compound; and
 - b) monitoring by any suitable means a binding interaction between said protein and said compound.
- 25 9. A pharmaceutical formulation comprising as an active ingredient TRAILLK-2 protein, associated with one or more pharmaceutically acceptable carriers, excipients, or diluents thereof.
10. TRAILLK-2 or an analog thereof for use in treating cancer in a patient in need thereof.

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 921 194 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
07.07.1999 Bulletin 1999/27

(51) Int Cl.⁶: **C12N 15/12, A61K 38/17,
G01N 33/68, C12N 1/21,
C07K 14/705, A61K 38/19,
C07K 14/52, C12N 15/19**

(43) Date of publication A2:
09.06.1999 Bulletin 1999/23

(21) Application number: **98309632.2**

(22) Date of filing: **25.11.1998**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(72) Inventor: **Song, Ho Yeong**
Carmel, Indiana 46032 (US)

(74) Representative: **Denholm, Anna Marie et al**
Eli Lilly and Company Limited,
Lilly Research Center,
Erl Wood Manor
Windlesham, Surrey GU20 6PH (GB)

(30) Priority: **26.11.1997 US 66577 P**
11.08.1998 US 96173 P

(71) Applicant: **ELI LILLY AND COMPANY**
Indianapolis, Indiana 46285 (US)

(54) **TRAILLK-2: a member of the TNF ligand family**

(57) The invention provides isolated nucleic acid compounds, proteins and fragments thereof, said proteins being related to the family of TNF ligands. Also provided are vectors and transformed heterologous host

cells for expressing the protein and a method for identifying compounds that bind and/or modulate the activity of said proteins, and a method for treating cancerous growths in a patient in need thereof.

EP 0 921 194 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 30 9632

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
P, X	WO 98 27114 A (SCHERING CORPORATION (US); GORMAN DANIEL M.) 25 June 1998 * abstract * * page 9 - page 15; table 1 * * page 39, line 18-32 * * page 47, line 6-20 * Seq.ID:3,4 * page 58 - page 61 * * page 67 - page 68; claims *	1-10	C12N15/12 A61K38/17 G01N33/68 C12N1/21 C07K14/705 A61K38/19 C07K14/52 C12N15/19
E	WO 99 12964 A (BIOGEN INC. (US); TSCHOPP JURG) 18 March 1999 * abstract * * page 5, line 16 - page 7, line 20 * * page 14, line 7-20 * * page 16, line 24 - page 17, line 5 * * page 23, line 10-19 * * page 28 - page 31; claims * Seq.ID:1,2 * page 32 * * figure 1 *	1-10	
P, X	WO 98 18921 A (HUMAN GENOME SCIENCES INC.; YU GUO-LIANG; EBNER REINHARD; NI JIAN) 7 May 1998 * abstract * * page 11, line 21 - page 12, line 11 * * page 21, line 7-15 * * page 41, line 19 - page 42, line 2 * * page 54 - page 57 * Seq.ID:1,2 * page 77 - page 80 * * page 90 - page 94; claims * * figures 1,2 *	1-10	C12N C07K A61K G01N
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 28 April 1999	Examiner Macchia, G
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document</p> <p>T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document</p>			

EPO FORM 1503 03/92 (P4/C21)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 30 9632

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
P,X	EP 0 869 180 A (SMITHKLINE BEECHAM CORPORATION (US); HURLE MARK R.; YOUNG PETER R.) 7 October 1998 * the whole document *	1-10	
E	WO 99 11791 A (UNIVERSITY OF WASHINGTON (US); CHAUDHARY PREET M.) 11 March 1999 * abstract * * page 30, line 3 - page 34, line 32 * * page 54, line 10-25 * * page 57, line 1 - page 59, line 12 * * page 129 - page 130; claims 33-38 * * figure 11A *	1-10	
E	WO 98 55620 A (REGENERON PHARMACEUTICALS (US); MASIAKOWSKI PIOTR; VALENZUELA DAVID) 10 December 1998 * the whole document *	1-10	
E	WO 98 55621 A (REGENERON PHARMACEUTICALS (US); MASIAKOWSKI PIOTR; VALENZUELA DAVID) 10 December 1998 * the whole document *	1-10	
X	Database EMBL, ID HSAA66695 Accession number AA166695 21 December 1996 98% identity with Seq.ID:1 nt.432-871 reverse orientation XP002101470 * the whole document *	3	
A	WO 97 33902 A (HUMAN GENOME SCIENCES INC. (US); NI JAN; YU GUO-LIANG; GENTZ REINER L.) 18 September 1997 * page 18, paragraph 3 * --- -/--		TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 28 April 1999	Examiner Macchia, G
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EP 0 921 194 A3 (P04001)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 30 9632

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A	GRUSS H.-J. AND DOWER S.K.: "Tumor necrosis factor ligand superfamily: involvement in the pathology of malignant lymphomas" BLOOD, vol. 85, no. 12, 15 June 1995, pages 3378-3404, XP002094502 ---		
A	AGGARWAL B.B. AND NATARAJAN K.: "Tumor necrosis factor: developments during the last decade" EUROPEAN CYTOKINE NETWORK, vol. 7, no. 2, April 1996 - June 1996, pages 93-124, XP002094503 ---		
A	GRUSS H.-J.: "Molecular, structural, and biological characteristics of the tumor necrosis factor ligand superfamily" INTERNATIONAL JOURNAL OF CLINICAL AND LABORATORY RESEARCH, vol. 26, no. 3, 1996, pages 143-159, XP002094504 -----		TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 28 April 1999	Examiner Macchia, G
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03/82 (P04/01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 30 9632

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

28-04-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9827114 A	25-06-1998	AU 5705898 A	15-07-1998
WO 9912964 A	18-03-1999	WO 9912965 A	18-03-1999
WO 9818921 A	07-05-1998	AU 7674596 A	22-05-1998
EP 0869180 A	07-10-1998	CA 2232743 A	02-10-1998
		JP 10323194 A	08-12-1998
WO 9911791 A	11-03-1999	NONE	
WO 9855620 A	10-12-1998	AU 7608898 A	21-12-1998
		AU 7713098 A	21-12-1998
		WO 9855621 A	10-12-1998
WO 9855621 A	10-12-1998	AU 7608898 A	21-12-1998
		AU 7713098 A	21-12-1998
		WO 9855620 A	10-12-1998
WO 9733902 A	18-09-1997	AU 5366596 A	01-10-1997
		EP 0897390 A	24-02-1999

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82